


ELECTROCHEMISTRY


(All questions are no calculator friendly.)

- 1) Balance each skeleton reaction, calculate $\varepsilon^\circ_{\text{cell}}$, and state whether or not the reaction is spontaneous.



(Answers: (a) Yes, $\varepsilon^\circ = +0.28 \text{ V}$, (b) No, $\varepsilon^\circ = -0.44 \text{ V}$, (c) No, $\varepsilon^\circ = -0.07 \text{ V}$)

- 2) Use the following half-reactions to write three spontaneous reactions, calculate ε° for each reaction, and rank the strength of the oxidizing and reducing agents:



(Answers: The three spontaneous reactions would be between:

- (1) $\text{Al}(s) + \text{N}_2\text{O}_4(\text{g})$**
- (2) $\text{Al}(s) + \text{SO}_4^{-2}(\text{aq})$**
- (3) $\text{SO}_4^{-2}(\text{aq}) + \text{NO}_2^{-1}(\text{aq})$**

Strongest to weakest reducing agent: $\text{Al}(s) > \text{NO}_2^{-1}(\text{aq}) > \text{SO}_3^{-2}(\text{aq})$

Strongest to weakest oxidizing agent: $\text{SO}_4^{-2}(\text{aq}) > \text{N}_2\text{O}_4(\text{g}) > \text{Al}^{+3}(\text{aq})$

- 3) A voltaic cell consists of a metal A/A⁺ electrode and a metal B/B⁺ electrode, with the A/A⁺ electrode negative. The initial [A⁺]/[B⁺] is such that $\varepsilon_{\text{cell}} > \varepsilon^\circ_{\text{cell}}$.

- a) How do [A⁺] and [B⁺] change as the cell operates?
- b) How does $\varepsilon_{\text{cell}}$ change as the cell operates?
- c) What is [A⁺]/[B⁺] when $\varepsilon_{\text{cell}} > \varepsilon^\circ_{\text{cell}}$? Explain.
- d) Is it possible for $\varepsilon^\circ_{\text{cell}}$ to be greater than $\varepsilon_{\text{cell}}$? Explain.

(Answers: (a) [A⁺] increases and [B⁺] decreases.

(b) $\varepsilon_{\text{cell}}$ decreases.

(c) [A⁺] < [B⁺]

(d) Yes, if [A⁺] > [B⁺]




- 4) A concentration cell consists of two Sn/Sn⁺² half-cells. The electrolyte in compartment A is 0.10 M Sn(NO₃)₂. The electrolyte in B is 1.00 M Sn(NO₃)₂. Which half-cell houses the cathode? What is the voltage of the cell?

(Answer: Compartment B houses the cathode and the cell voltage is +29.6 mV.)




5) A voltaic cell with Mn/Mn²⁺ and Cd/Cd²⁺ half-cells has the following starting concentrations: [Mn²⁺] = 0.010 M and [Cd²⁺] = 0.100 M.

- What is the initial ϵ_{cell} ? **(Answer: $\epsilon_{\text{cell}} = +0.81 \text{ V}$)**
- What is ϵ_{cell} when [Mn²⁺] = 0.055 M? **(Answer: $\epsilon_{\text{cell}} = +0.78 \text{ V}$)**
- Calculate K_{eq} and the concentrations of the ions at equilibrium.
(Answers: $K_{\text{eq}} \approx 1 \times 10^{26}$ and [Mn²⁺] = 0.110 M and [Cd²⁺] = 1 x 10⁻²⁷ M)

6) The following reactions occur at 25°C with all soluble substances present in 1 M concentrations:



From this information alone, predict whether the following reactions will occur under similar conditions:



YOUR ANSWERS MUST BE RATIONALIZED BY INDICATING A RELATIVE SCALE OF EMF FOR THE FOUR ELEMENTS IN QUESTION.

(Answers: (a) YES (b) NO (c) YES)

7) Explain why copper is oxidized by nitric acid and not by hydrochloric acid.


(Answer: Copper cannot be oxidized by HCl because for the reaction $\text{Cu} + 2\text{H}^+ \longrightarrow \text{H}_2 + \text{Cu}^{2+}$ the $\epsilon^\circ_{\text{cell}}$ is less than zero. However for the reactions $\text{Cu} + \text{HNO}_3 \longrightarrow \text{Cu}^{2+} + \text{NO}_2$ or NO the $\epsilon^\circ_{\text{cell}}$ is greater than zero.)

8) The corrosion of iron involves oxidation of the metal and the subsequent formation of the oxide. Which metal, zinc or nickel, will provide the better protection against corrosion if coated in a thin layer on the iron?

(Answer: Zinc would be better than nickel because it has a more positive value for its oxidation potential than does iron, and thus would be oxidized more readily than iron. However if it were coated with nickel, the iron would be oxidized before the nickel and thus would not protect the iron.)

- 9) A solution contains the following metal ions: $[\text{Au}^{3+}] = 1.0 \times 10^{-6} \text{ M}$, $[\text{Fe}^{3+}] = 1.0 \text{ M}$ and $[\text{Ni}^{2+}] = 1.0 \times 10^{-4} \text{ M}$. What metal will plate out first if this solution is electrolyzed? (HINT: Use the Nernst equation to calculate ϵ for each half-reaction.)
(Answer: For $[\text{Au}^{3+}] = 1.0 \times 10^{-6} \text{ M}$ the $\epsilon_{\text{half-cell}} = +1.38 \text{ V}$
For $[\text{Fe}^{3+}] = 1.0 \text{ M}$ the $\epsilon_{\text{half-cell}} = -0.037 \text{ V}$
For $[\text{Ni}^{2+}] = 1.0 \times 10^{-4} \text{ M}$ the $\epsilon_{\text{half-cell}} = -0.37 \text{ V}$
Since the $\epsilon_{\text{half-cell}}$ for the Au^{3+}/Au is the most positive it will be the first metal to plate out of the solution.)

- 10) A voltaic cell employs the following two half-reactions:



Write the overall cell reaction and calculate ϵ_{cell} if the pH = 4.00 and all other substances are at standard conditions. **(Answer: $\epsilon_{\text{cell}} = +0.52 \text{ V}$)**

- 11) Hydrolysis of molten MgCl_2 is the final production step in the isolation of magnesium from seawater by the Dow process. Assuming that 48.6 g of Mg forms:

- How many moles of electrons are required?
- How many coulombs are required?
- How many amps will produce this amount in 11 hours?

(Answers: (a) 4.0 mol e^{-1} (b) $4 \times 9.65 \times 10^4 \text{ C} \approx 4 \times 10^5 \text{ C}$ (c) about 10 A)

- 12) Zinc plating (galvanizing) is an important means of corrosion protection. Although the process is done customarily by dipping the object into molten zinc, the metal can also be electroplated from aqueous solutions. How many grams of zinc can be deposited on a steel tank from a ZnSO_4 solution when a 0.965 A current flows for 2.3 days (approx. 2.0×10^5 seconds)?


(Answer: 1 mole zinc \approx 65 g zinc)

- 13) Car manufacturers are developing engines that use H_2 as fuel. In Iceland, Sweden, and other parts of Scandinavia, where hydroelectric plants produce inexpensive electric power, the H_2 can be made industrially by the electrolysis of water.

- How many coulombs are needed to produce $2.5 \times 10^6 \text{ L}$ of H_2 gas at 10.0 atm and 25°C ? (Assume the ideal gas law applies.)
- If the coulombs are supplied at 1.5 V, how many joules are produced?
- If the combustion of oil yields $4.0 \times 10^4 \text{ kJ/kg}$, what mass of oil must be burned to yield the number of joules in part (b)?

(Answers: (a) $2.0 \times 10^{11} \text{ C}$ (b) $3.0 \times 10^{11} \text{ J}$ (c) $7.5 \times 10^3 \text{ kg oil}$)


14) Electrodes used in electrocardiography are disposable, and many incorporate silver. The metal is deposited in a thin layer on a small plastic "button", and then some is converted to AgCl:




- If the surface area of the button is 2.0 cm^2 and the thickness of the silver layer is $7.0 \times 10^{-6} \text{ m}$, calculate the volume (in cm^3) of Ag used in one electrode.
- The density of silver metal is 10.5 g/cm^3 . How many grams of silver are used per electrode?
- If the silver is plated on the button from an Ag^+ solution with a current of 10.0 mA , how many minutes does the plating take?
- If bulk silver costs \$5.50 per troy ounce (31.10 g), what is the cost (in cents) of the silver in one disposable electrode?

(Answers: (a) $1.4 \times 10^{-3} \text{ cm}^3$ (b) $1.5 \times 10^{-2} \text{ g Ag}$ (c) about 25 min (d) about 0.3 cents)

15) Given the following standard reduction potentials at 25°C



- Calculate the value of the equilibrium constant at 25°C for the reaction:



(Answer: $K_{\text{eq}} = 1 \times 10^{13}$)

- Write the conventional cell notation for the cell you have sketched.

(Answer: $\text{Ag(s)} | \text{S}_2\text{O}_3^{-2}(\text{aq}), \text{cations}(\text{aq}), \text{Ag(S}_2\text{O}_3)_2^{-3}(\text{aq}) || \text{Ag}^+(\text{aq}), \text{anions}(\text{aq}) | \text{Ag(s)}$)

- Sketch the voltaic cell in which the above chemical equation is the overall reaction. Your sketch should clearly indicate the following: anode, cathode, electrode signs, electron movement as well as the reactions taking place both the anode and cathode.

Answer:

